

Infliximab (Remicade or other infliximab product as required by patient's health plan)

Fax to 505-420-4848 or Email to refer@optimuminfusion.com



PATIENT INFORMATION

Referral Status: New Referral Updated Order Order Renewal

Date: _____ Patient Name: _____ DOB: _____

ICD-10 code (required): _____ ICD-10 description: _____

NKDA Allergies: _____ Weight (lbs/kg): _____ Height: _____

Patient Status: New to Therapy Continuing Therapy Last Treatment Date: _____ Next Due Date: _____

PROVIDER INFORMATION

Referral Coordinator Name: _____ Referral Coordinator Email: _____

Ordering Provider: _____ Provider NPI: _____

Referring Practice Name: _____ Phone: _____ Fax: _____

Practice Address: _____ City: _____ State: _____ Zip Code: _____

NURSING

- TB status & date (list results here and attach clinicals)

- Hepatitis B status & date (list results here and attach clinicals)

- Provide nursing care per Optimum Infusion Nursing Procedures, including reaction management and post-procedure observation

LABORATORY ORDERS

- CBC at each dose every _____
- CMP at each dose every _____
- CRP at each dose every _____
- Other: _____

PRE-MEDICATION ORDERS

- acetaminophen (Tylenol) 500mg / 650mg / 1000mg PO
 - cetirizine (Zyrtec) 10mg PO
 - loratadine (Claritin) 10mg PO
 - diphenhydramine (Benadryl) 25mg / 50mg PO / IV
 - methylprednisolone (Solu-Medrol) 40mg / 125mg IV
 - hydrocortisone (Solu-Cortef) 100mg IV
 - Other: _____
- Dose: _____ Route: _____
- Frequency: _____

SPECIAL INSTRUCTIONS

THERAPY ADMINISTRATION

Infliximab (Remicade) or other infliximab product (as required by patient's health plan)

NOTE: (Infliximab products include: Remicade, Unbranded Infliximab, Avsola, Inflectra, and Renflexis)

Dose: 3mg/kg 5mg/kg 7.5mg/kg 10mg/kg

Other: _____

Round up to nearest 100mg **OR** Give exact dose

Frequency: induction: week 0, 2, 6, and then every 8 weeks /

maintenance: every 8 weeks / other: _____

Infusion rate: Select one below. Patients who tolerate induction and the initial maintenance infusion without severe reaction will be eligible for 1 hour infusion

Infuse over 2 hours (standard rate)

Infuse over 1 hour (when patient eligible)

Flush with 0.9% sodium chloride at infusion completion

Refills: Zero / for 12 months / _____
(if not indicated order will expire one year from date signed)

*Perform test for latent TB; if positive, start treatment for TB prior to starting treatment. Monitor all patients for active TB during treatment, even if initial latent TB test is negative. *Patients should be tested for HBV infection before initiating TNF blocker therapy, including REMICADE. For patients who test positive for hepatitis B surface antigen, consultation with a physician with expertise in the treatment of hepatitis B is recommended.

Provider Name (Print)

Provider Signature

Date

SUBMIT ORDER FORM TO OPTIMUM INFUSION:

FAX: 505-420-4848

EMAIL: refer@optimuminfusion.com